

# Property Library for Isohexane

## LibC6H14

Prof. Hans-Joachim Kretzschmar  
Dr. Sebastian Herrmann  
Dr. Matthias Kunick  
Ines Jaehne  
Tobias Goepfert

## Property Functions

Functional Dependence	Function Name	Call from Fortran Program	Property or Function	Unit of the Result
$a = f(p, t, x)$	a_ptx_C6H14	A_PTX_C6H14(P,T,X)	Thermal diffusivity	m <sup>2</sup> /s
$c_p = f(p, t, x)$	cp_ptx_C6H14	CP_PTX_C6H14(P,T,X)	Specific isobaric heat capacity	kJ/(kg K)
$c_v = f(p, t, x)$	cv_ptx_C6H14	CV_PTX_C6H14(P,T,X)	Specific isochoric heat capacity	kJ/(kg K)
$\eta = f(p, t, x)$	eta_ptx_C6H14	ETA_PTX_C6H14(P,T,X)	Dynamic viscosity	Pa . s
$h = f(p, t, x)$	h_ptx_C6H14	H_PTX_C6H14(P,T,X)	Specific enthalpy	kJ/kg
$\kappa = f(p, t, x)$	ka_ptx_C6H14	KA_PTX_C6H14(P,T,X)	Isentropic exponent	-
$\lambda = f(p, t, x)$	lam_ptx_C6H14	LAM_PTX_C6H14(P,T,X)	Thermal conductivity	W/(m . K)
$\nu = f(p, t, x)$	ny_ptx_C6H14	NY_PTX_C6H14(P,T,X)	Kinematic viscosity	m <sup>2</sup> /s
$Pr = f(p, t, x)$	pr_ptx_C6H14	PR_PTX_C6H14(P,T,X)	<i>Prandtl</i> -number	-
$p_s = f(t)$	ps_t_C6H14	PS_T_C6H14(T)	Vapor pressure from temperature	bar
$\rho = f(p, t, x)$	rho_ptx_C6H14	RHO_PTX_C6H14(P,T,X)	Density	kg/m <sup>3</sup>
$s = f(p, t, x)$	s_ptx_C6H14	S_PTX_C6H14(P,T,X)	Specific entropy	kJ/(kg K)
$\sigma = f(t)$	sigma_t_C6H14	SIGMA_T_C6H14(T)	Surface tension from temperature	N/m
$t = f(p, h)$	t_ph_C6H14	T_PH_C6H14(P,H)	Backward function: Temperature from pressure and enthalpy	°C
$t = f(p, s)$	t_ps_C6H14	T_PS_C6H14(P,S)	Backward function: Temperature from pressure and entropy	°C
$t_s = f(p)$	ts_p_C6H14	TS_P_C6H14(P)	Saturation temperature from pressure	°C
$u = f(p, t, x)$	u_ptx_C6H14	U_PTX_C6H14(P,T,X)	Specific internal energy	kJ/kg
$v = f(p, t, x)$	v_ptx_C6H14	V_PTX_C6H14(P,T,X)	Specific volume	m <sup>3</sup> /kg
$w = f(p, t, x)$	w_ptx_C6H14	W_PTX_C6H14(P,T,X)	Isentropic speed of sound	m/s

Functional Dependence	Function Name	Call from Fortran Program	Property or Function	Unit of the Result
$x = f(p, h)$	x_ph_C6H14	X_PH_C6H14(P,H)	Backward function: Vapor fraction from pressure and enthalpy	kg/kg
$x = f(p, s)$	x_ps_C6H14	X_PS_C6H14(P,S)	Backward function: Vapor fraction from pressure and entropy	kg/kg

**Units:**

- $t$  in °C
- $p$  in bar
- $x$  in (kg saturated steam)/(kg wet steam)

### Range of validity

for transport properties ( $a$ ,  $\eta$ ,  $\lambda$ ,  $\nu$ ,  $Pr$ ):

Temperature range: from - 153.55 °C to 276.85 °C  
Pressure range: from  $7.6739 \times 10^{-11}$  bar to 1000 bar

for other properties:

Temperature range: from - 153.55 °C to 276.85 °C  
Pressure range: from  $7.6739 \times 10^{-11}$  bar to  $1 \times 10^4$  bar

### Reference state

$h = 0$  kJ/kg and  $s = 0$  kJ/(kg K) at  $p = 1,01325$  bar on the saturated liquid line ( $x = 0$ )

## Details on the vapor fraction $x$

The wet steam region is calculated automatically by the subprograms. For this purpose the following fixed details on the vapor fraction  $x$  are to be considered:

### Single-phase region

If the state point to be calculated is located in the single-phase region (liquid or superheated steam)  $x = -1$  must be entered as a pro-forma value.

### Wet-steam region

If the state point to be calculated is located in the wet steam region, a value for  $x$  between 0 and 1 ( $x = 0$  for saturated liquid,  $x = 1$  for saturated steam) must be entered. In this case, the backward functions result in the appropriate value between 0 and 1 for  $x$ . When calculating wet steam either the given value for  $t$  and  $p = -1000$  or the given value for  $p$  and  $t = -1000$  and in both cases the value for  $x$  between 0 and 1 must be entered.

If  $p$  and  $t$  and  $x$  are entered as given values, the program considers  $p$  and  $t$  to be appropriate to represent the vapor pressure curve. If this is not the case the calculation for the property of the chosen function results in  $-1$ .

Wet steam region: Temperature ranges from  $t_{\min} = -153.55 \text{ °C}$  to  $t_c = 224.55 \text{ °C}$   
Pressure ranges from  $p_{\min} = 7.6739 \times 10^{-11} \text{ bar}$  to  $p_c = 30.426 \text{ bar}$

### **Note:**

*If the input values are located outside the range of validity, the calculated function will always result in  $-1000$ . More exact details on every function and its corresponding range of validity can be found in the enclosed program documentation in chapter 3.*